

## Catalogue of American Amphibians and Reptiles.

PICKWELL, GEORGE V., AND WENDY A. CULOTTA. 1980. *Pelamis*, *P. platurus*.

*Pelamis* Daudin

## Pelagic or yellow-bellied sea snake

*Hydrus* Schneider, 1799:233 (in part). Type-species, *Hydrus hydrus* (Pallas) 1771 (= *Hydrus caspius* Schneider, 1799, = *Coluber hydrus* Pallas, 1771), by original designation. See NOMENCLATURAL HISTORY.

*Hydrophis* Latreille, 1801:193 (in part). No type-species designated. See NOMENCLATURAL HISTORY.

*Pelamis* Daudin, 1803:357 (in part). Type-species, *Pelamis bicolor* (Schneider) (= *Anguis platyura* Linnaeus, 1766), by designation of Gray, 1825.

*Pelamys* Wagler, 1830:166. Emendation of the Greek spelling to an equivalent form.

*Pelamydrus* Stejneger, 1911:111. See NOMENCLATURAL HISTORY.

• CONTENT. The single species recognized is *Pelamis platurus* (Linnaeus), 1766.

• DEFINITION, DIAGNOSIS, DESCRIPTIONS, ILLUSTRATIONS, DISTRIBUTION, PERTINENT LITERATURE. See species account.

• FOSSIL RECORD. No fossils of this genus are known.

• NOMENCLATURAL HISTORY. The generic name for this snake has undergone a long and convoluted history. Schneider (1799) assigned the species *bicolor* to his genus *Hydrus*, which also contained Pallas' *Coluber hydrus* (= *Hydrus caspius* Schneider), a freshwater snake that was ultimately made the type for *Hydrus* (Stiles, 1910). Latreille (1801), evidently considering *Hydrus* too similar to *Hydra*, substituted *Hydrophis*. But shortly thereafter, Daudin (1803), considering *Hydrophis* an inappropriate name for marine snakes, substituted *Pelamis*. Why this appealed to Daudin as a name for a genus of serpents is unclear. Gray (1825) recognized the uniqueness of this animal when he designated *Anguis platyura* Linnaeus the type of the monotypic genus *Pelamis*. Much later, Boulenger (1890), for unknown reasons, reinstated *Hydrus* as a monotypic genus with *A. platyura* Linnaeus as the type. Following the decision of the International Commission on Zoological Nomenclature (Stiles, 1910) making *Hydrus caspius* the type of Schneider's genus *Hydrus*, Stejneger (1911) substituted *Pelamydrus* for *Hydrus* (as employed by Boulenger, 1890). Ultimately, M. Smith (1926) reinstated *Pelamis* on the grounds that it was the oldest available name. Recent authorities (McDowell, 1972; Burger and Natsuno, 1974; Cogger, 1975a, b) have continued use of the monotypic genus *Pelamis*. For discussions pertaining to various nomenclatural changes, see Stejneger (1911), Smith (1926), Smith and Taylor (1945), and Loveridge (1957).

• ETYMOLOGY. *Pelamis* is an apparently obsolete Greek noun (masculine) referring to a young tuna.

*Pelamis platurus* (Linnaeus)  
Pelagic or yellow-bellied sea snake

*Anguis platyura* Linnaeus, 1766:391. Type-locality and holotype unknown.

*Anguis platurus* Linnaeus, 1788:1122. Type-locality, "Pine-isle." Holotype unknown.

*Hydrus bicolor* Schneider, 1799:242. Type-locality, "Otaheitee." Holotype unknown. Account based on descriptions by Seba (1735), Vosmaer (1774), Linnaeus (1788) and Russell (1796). See NOMENCLATURAL HISTORY in genus account.

*Hydrophis platyura*: Latreille, 1801:197. New combination. See NOMENCLATURAL HISTORY in genus account.

*Pelamis bicolor*: Daudin, 1803:366. New combination. See NOMENCLATURAL HISTORY in genus account.

*Hydrophis pelamis* Schlegel in Temminck and Schlegel, 1838:90. New combination. This name appeared in Siebold's Fauna Japonica. For the correct dates of the sections from Reptilia consult Sherborn and Jentink (1895).

*Hydrophis (Pelamis) bicolor*: Fischer, 1856:61. Substitute name for *H. pelamis* Schlegel giving the trivial name subgeneric status.

*Pelamys bicolor*: Fitzinger, 1861:409. Emendation of *P. bicolor* Daudin based on the revised generic spelling of Wagler (1830).

*Pelamis platurus*: Stoliczka, 1870:214. New combination.


*Hydrus platurus*: Boulenger, 1890:397. Substitute name for *P. platurus* Stoliczka.

*Pelamydrus platurus*: Stejneger, 1911:111. Substitute name for *H. platurus* Boulenger.


*Pelamis platyura*: Deraniyagala, 1955:79. Emendation of species ending to conform with the original Linnaean designation.

• CONTENT. The species is monotypic.

• DEFINITION AND DIAGNOSIS. This species is venomous and live-bearing. Brood size varies generally from 2 to 8. Adults seldom exceed 75 cm total length. The largest specimens are from eastern Australian waters (maximum length 113 cm). The body is highly compressed except in gravid individuals; the ventrals small, approximately 260 to greater than 400 in long specimens, usually divided, with the groove forming a straight mid-ventral sulcus, the ventrals scarcely wider than adjacent scales. The tail is flattened like an oar; body scales are small, juxtaposed and more or less hexagonal or quadrangular, in usually 49 to 67 rows on the thickest part of the body, with about three transverse rows per vertebra at midbody, occasionally bearing tubercles in adults. The head is elongate, narrow and flattened; fangs are small, 1 to 2 mm long, about one-half of the distance from the rostrum to mid-orbit and separated from the solid maxillary teeth by a con-



MAP 1. Coarsely stippled area represents the worldwide distribution of *Pelamis* based on collected specimens and reliable sightings. Densely shaded regions are areas of probable occasional occurrence depending on ocean current conditions, sea surface temperature and season.



MAP 2. Shaded area indicates the region of common occurrence of *Pelamis* either as breeding populations or frequent strays. The distance offshore within which the snake occurs in quantity is highly variable. Circles represent remote areas where specimens have been identified; question marks indicate general localities of uncertain sightings reported in the literature, but with no exact location data or voucher specimens.

spicuous diastema. Maxillary teeth are usually 7 to 10, palatine teeth, 6 or 7. Head shields are enlarged, smooth and without tubercles or spines, nasals are in contact on the dorsal surface of the snout. There are no internasals; the nares are superior and valved. The hemipenis is narrow and only slightly bilobate with papillae at its tip; the sulcus spermaticus is forked near the tip, and the organ is covered over most of its length by small spines, each about as long as the width of the caudal scales.

Color patterns involve a sharp lateral demarcation between a black or dark brown dorsum and a bright yellow or brownish venter; in the latter case the two zones are often separated by a yellow line of varying width. The tail is generally spotted or barred with black or dark brown on yellow, no two individuals ever having exactly the same pattern. There is no evidence at present for significant ontogenetic variation in pattern or color intensity. Other sea snakes possessing black or dark brown and yellow coloration are invariably banded. There are numerous variations on the basic bicolor or tricolor scheme in *Pelamis* involving undulating patterns, additional dark lines, bars or spots, to the extreme and rare cases of specimens bearing bars the entire length of the body or lacking dark pigmentation altogether.

• **DESCRIPTIONS.** There is no definitive taxonomic or morphological treatment of *Pelamis*, but Wall (1921) and Smith (1926) gave useful brief accounts. Valuable early comparative studies include: buccal glands and teeth (West, 1895); olfactory glands (Katheriner, 1900); visceral anatomy (Beddard, 1904); cranial and tooth structure, cephalic musculature and glands (Phisalix, 1912, 1914); cephalic muscles (Haas, 1930). Contemporary comparative anatomical works include: venom glands (Rosenberg, 1967); hyoid (Langebartel, 1968); oral sensory systems (Burns, 1969a, b); cephalic glands (Burns and Pickwell, 1972); jaw muscles and cranial structure (McDowell, 1972); retinal morphology (Hibbard and LaVergne, 1972); eyes and other sense organs (Hibbard, 1975); lung morphology (Graham et al., 1975; Heatwole and Seymour, 1975); salt glands and integument (Dunson, 1975a, 1976); dermal scale-vertebral relationships (Voris, 1975); costocutaneous muscles (Voris and Jayne, 1976). Data on the numerous color and pattern variations were provided by Barbour (1912), Deraniyagala

(1960), Visser (1967, includes observations on brood size), Kropach (1972a) and Bolaños et al. (1974).

• **ILLUSTRATIONS.** Black and white photographs of *Pelamis* showing pattern variations appeared in Picado (1976), Shaw (1961), Paulson (1967), Rubinoff and Kropach (1970) and Cogger (1975b). Photos of *Pelamis* undertaking knotting maneuvers appeared in Pickwell (1971). Kropach (1971a) depicted an all-yellow specimen. An excellent close-up of the integument showing scale detail was given in Dunson (1975a). Color photos of value were presented by Kuntz (1963), Halstead (1970), Dunson (1971), Pickwell (1972) and Tu (1976).

• **DISTRIBUTION.** *Pelamis* is the most widely distributed of all snake species, but is not known to occur in the Atlantic. Shuntov (1963) and Marinkelle (1966) reported sightings of *Pelamis* in the Caribbean, but without verification by collected specimens. The literature containing distributional data on *Pelamis* is massive. Many of the major features of its distribution were known one hundred years ago and were well documented in Günther (1864), Strauch (1873) and Peters and Doria (1878). Also see Boulenger (1896) and Werner (1900) for extensive distribution lists and Bourret (1934) for an interesting attempt to correlate Smith's (1926) varieties with geographic regions. Minton (1975) has provided the most recent comprehensive general distribution lists.

• **NOMENCLATURE HISTORY.** Gray's (1842) *Pelamis ornata* has frequently been cited in the synonymy of *P. platurus*. But this designation was a substitute name for a variety of *Hydrus bicolor* described by Schlegel (in Temminck and Schlegel, 1838) and cannot be considered in the main line of *P. platurus* synonymy. Furthermore, Gray listed this variation immediately following his description of *P. bicolor* Daudin. Smith (1920) was of the opinion that Barbour's (1912) subspecies, *ornatus*, was insufficiently substantiated on geographic grounds alone, since other varieties occurred in the same area. But see Taylor (1950) for an opposing view. Wall (1919, 1921) was unfortunately inconsistent in his varietal designations and evidently established new synonymy lists in each new publication dealing with *P. platurus*.

Rafinesque (1817) erected the species *luteus* for an all-yellow snake in his genus, *Ophionectes*. This species was designated the type of the genus by Loveridge (1957), who believed that it and "some or all" of the remaining nine new species in the genus were but variants of *P. platurus*. This seems most unlikely since among these were included species that were entirely gray, green or bluish, with a red head, or variously spotted or banded with blue, white, red or green. For many of these it is uncertain that Rafinesque had actually seen any snake specimens at all. The first reliable mention of an all-yellow *Pelamis* was that of Voris et al. (1970) followed by the definitive description by Kropach (1971a) and corroborative observations by Tu (1976). It is not clear from the literature that either Rafinesque or Loveridge were aware of this rare color phase.

• **PERTINENT LITERATURE.** The definitive monograph of *Pelamis* general biology is that of Kropach (1972a). Also see Kropach (1975) for a shortened version. Briefer general treatments based on field and laboratory studies appeared by Dunson (1971) and Pickwell (1972). Viewpoints on the significance of *Pelamis* as a potential colonizer of the Caribbean and Atlantic via a possible Middle American sea level canal were given by Dunson (1971, 1975b) and opposing views by Kropach (1972b), Pickwell (1972) and Pickwell et al. (1973). The significance of ocean slicks to this snake's feeding and general life history was documented by Dunson and Ehlert (1971) and Kropach (1971b, 1972a). Prey items used by *Pelamis* were listed by Clark (1942), Klawe (1964), Visser (1967), Voris (1969, 1971) and Kropach (1971b, 1972a). Feeding behavior was described by Klauber (1935), Shaw (1962), Zeiller (1969), Kropach (1972a) and Pickwell (1972). The evident lack of predators and possible aposematic use of the snake's color and pattern were discussed by Rubinoff and Kropach (1970) and Kropach (1972a). The effects of ocean surface temperature on the survival and distribution of *Pelamis* were studied by Graham et al. (1971), Dunson and Ehlert (1971), Graham (1974a) and Hecht et al. (1974). Aspects of respiration and diving physiology were reported by Graham (1974b), Graham et al. (1975), Heatwole and Seymour (1975) and Heatwole (1978). The osmoregulatory and salt excretion capacities of *Pelamis* were well documented in the work of Dunson and colleagues (Dunson, 1975a, 1976; Dunson and Robinson, 1976).

• **ETYMOLOGY.** *Platurus* is a compound of the Greek *platys* (or *platos*) meaning "flat," and *oura*, "tail."

## LITERATURE CITED

- Barbour, Thomas. 1912. A contribution to the zoogeography of the East Indian islands. Mem. Mus. Comp. Zool. 44(1):1-203.
- Beddard, Frank E. 1904. Contributions to the knowledge of the visceral anatomy of the pelagic serpents *Hydrus platyurus* and *Platyurus colubrinus*. Proc. Zool. Soc. London 2:147-154.
- Bolaños, Róger, Alvaro Flores, Richard T. Taylor, and Luis Cerdas. 1974. Color patterns and venom characteristics in *Pelamis platurus*. Copeia 1974(4):909-912.
- Boulenger, George A. 1890. Reptilia and Batrachia. In W. T. Blanford (ed.), The fauna of British India, including Ceylon and Burma. Taylor and Francis, London. xviii + 541 p.
- 1896. Catalogue of the snakes in the British Museum (Natural History). Vol. 3. London. xiv + 727 p.
- Bourret, René. 1934. Les serpents marins de l'Indochine Française. Inst. Océanogr. Indochine, note 25. 69 p.
- Burger, W. Leslie, and Tetsuya Natsuno. 1974. A new genus for the Arafura smooth seasnake and redefinitions of other seasnake genera. Snake 6(2):61-75.
- Burns, Barry. 1969a. Oral sensory papillae in sea snakes. Copeia 1969(3):617-619.
- 1969b. An oral sensory system in sea serpents (Hydrophiidae); histological and morphological considerations. M.A. Thesis. Univ. Waterloo. vii + 22 p.
- , and George V. Pickwell. 1972. Cephalic glands in sea snakes (*Pelamis*, *Hydrophis* and *Laticauda*). Copeia 1972(3):547-559.
- Clark, Herbert C. 1942. Venomous snakes. Some Central American records. Incidence of snake-bite accidents. Amer. J. Trop. Med. 22(1):37-49.
- Cogger, Harold G. 1975a. Reptiles and amphibians of Australia. A. H. and A. W. Reed, Sydney, Australia. 584 p.
- 1975b. Sea snakes of Australia and New Guinea, p. 59-139. In William A. Dunson (ed.), The biology of sea snakes. University Park Press, Baltimore, Maryland. x + 530 p.
- Daudin, François M. 1803. Histoire naturelle, générale et particulière des reptiles. Tome septième. Dufart, Paris. 436 p.
- Deraniyagala, P. E. P. 1955. A colored atlas of some vertebrates from Ceylon. Vol. 3. Serpentine Reptilia. Ceylon Nat. Mus. Publ. xviii + 121 p.
- 1960. Some southern Temperate Zone snakes, birds and whales that enter the Ceylon area. Spolia Zeylanica 29(1):79-85.
- Dunson, William A. 1971. The sea snakes are coming. Natur. Hist. 80(9):52-61.
- 1975a. Salt and water balance in sea snakes, p. 329-353. In William A. Dunson (ed.), The biology of sea snakes. University Park Press, Baltimore, Maryland. x + 530 p.
- 1975b. Sea snakes and the sea level canal controversy. Op. cit., p. 517-524.
- 1976. Salt glands in reptiles, p. 413-445. In Carl Gans and William R. Dawson (eds.), Biology of the Reptilia, vol. 5. Physiology A. Academic Press, London, New York and San Francisco. xv + 556 p.
- , and Gary W. Ehlert. 1971. Effects of temperature, salinity and surface water flow on distribution of the sea snake *Pelamis*. Limnol. Oceanogr. 16(6):845-853.
- , and Gerald D. Robinson. 1976. Sea snake skin: permeable to water but not to sodium. J. Comp. Physiol. 108:303-311.
- Fischer, Johann G. 1856. Die Familie der Seeschlangen. Abhandl. Geb. Naturwiss. Hamburg. 3:1-78.
- Fitzinger, Leopold J. 1861. Die Ausbeute der österreichischen Naturforscher an Säugethieren und Reptilien während der Weltumsegelung Sr. Majestät Fregatte Novara. Sitzungsber. Akad. Wiss. Math.-Naturwiss. Klasse, Vienna. 42(25):383-416.
- Graham, Jeffrey B. 1974a. Body temperatures of the sea snake *Pelamis platurus*. Copeia 1974(2):531-533.
- 1974b. Aquatic respiration in the sea snake *Pelamis platurus*. Resp. Physiol. 21(1):1-7.
- , I. Rubinoff, and M. K. Hecht. 1971. Temperature physiology of the sea snake *Pelamis platurus*: An index of its colonization potential in the Atlantic Ocean. Proc. Nat. Acad. Sci. USA 68(6):1360-1363.
- , John H. Gee, and Fred S. Robison. 1975. Hydrostatic and gas exchange functions of the lung of the sea snake *Pelamis platurus*. Comp. Biochem. Physiol. 50A:477-482.
- Gray, John E. 1825. A synopsis of the genera of reptiles and Amphibia, with a description of some new species. Ann. Phil. new ser. 10:193-217.
- 1842. Monographic synopsis of the water snakes, or the family Hydriidae. Zool. Misc. 1842:59-68.
- Günther, Albert C. L. G. 1864. The reptiles of British India. Ray Soc. Publ. (35). xxvii + 452 p.
- Haas, Georg. 1930. Über die Schädelmechanik und die Kiefermuskulatur einiger Proteroglypha. Zool. Jahrb. Abt. Anat. Ont. Tiere 52(2):347-404.
- Halstead, Bruce W. 1970. Poisonous and venomous marine animals of the world. Vol. 3. Govt. Print. Off., Washington, D.C. xxv + 1006 p.
- Heatwole, Harold. 1978. Adaptations of marine snakes. Amer. Sci. 66(5):594-604.
- , and Roger Seymour. 1975. Diving physiology, p. 289-327. In William A. Dunson (ed.), The biology of sea snakes. University Park Press, Baltimore, Maryland. x + 530 p.
- Hecht, Max K., Chaim Kropach, and Bessie M. Hecht. 1974. Distribution of the yellow-bellied sea snake, *Pelamis platurus*, and its significance in relation to the fossil record. Herpetologica 30(4):387-396.
- Hibbard, Emerson. 1975. Eyes and other sense organs of sea snakes, p. 355-382. In William A. Dunson (ed.), The biology of sea snakes. University Park Press, Baltimore, Maryland. x + 530 p.
- and J. LaVergne. 1972. Morphology of the retina of the sea snake, *Pelamis platurus*. J. Anat. 112(1):125-136.
- Kathariner, Ludwig. 1900. Die Nase der im Wasser lebenden Schlangen als Luftweg und Geruchsorgan. Zool. Jahrb. Abt. Syst. Oekol. Geogr. Tiere 13:415-442.
- Klauber, Laurence M. 1935. The feeding habits of a sea snake. Copeia 1935(4):182.
- Klawe, Witold L. 1964. Food of the black and yellow sea snake, *Pelamis platurus* from Ecuadorian waters. Copeia 1964(4):712-713.
- Kropach, Chaim. 1971a. Another color variety of the sea snake *Pelamis platurus* from Panama Bay. Herpetologica 27(3):326-327.
- 1971b. Sea snake (*Pelamis platurus*) aggregations on slicks in Panama. Ibid. 27(2):131-135.
- 1972a. A field study of the sea snake *Pelamis platurus* (Linnaeus) in the Gulf of Panama. Ph.D. Thesis, City Univ. New York. x + 200 p.
- 1972b. *Pelamis platurus* as a potential colonizer of the Caribbean Sea. Bull. Biol. Soc. Washington. 2:267-269.
- 1975. The yellow-bellied sea snake, *Pelamis* in the eastern Pacific, p. 185-213. In William A. Dunson (ed.), The biology of sea snakes. University Park Press, Baltimore, Maryland. x + 530 p.
- Kuntz, Robert E. 1963. Snakes of Taiwan. Quart. J. Taiwan Mus. 16(1-2):1-79.
- Langebartel, David A. 1968. The hyoid and its associated muscles in snakes. Illinois Biol. Monogr. (38):1-156.
- Latreille, Pierre A. 1801. In C. S. Sonnini and P. A. Latreille, Histoire naturelle des reptiles. Détéville, Paris. Vol. 4, second part. Serpens. 410 p.
- Linnaeus, Carolus. 1766. Systema naturae per regna tria naturae . . . editio duodecima, reformata. Vol. 1. Laurentius Salvius, Holmiae. 532 p.
- 1788. Systema naturae. Edition 13. J. F. Gmelin (ed.). G. E. Beer, Lipsiae. Vol. 1, pt. 3: 1031-1516.
- Loveridge, Arthur. 1957. Check list of the reptiles and amphibians of East Africa (Uganda; Kenya; Tanganyika; Zanzibar). Bull. Mus. Comp. Zool. 117(2):153-362.
- McDowell, Samuel B. 1972. The genera of sea-snakes of the *Hydrophis* group (Serpentes: Elapidae). Trans. Zool. Soc. London 32:189-247.
- Marinkelle, C. J. 1966. Accidents by venomous animals in Colombia. Industrial Med. Surg. 1966:988-992.
- Minton, Sherman A. 1975. Geographic distribution of sea snakes, p. 21-31. In William A. Dunson (ed.), The biology of sea snakes. University Park Press, Baltimore, Maryland. x + 530 p.
- Pallas, Peter S. 1771. Reise durch verschiedene Provinzen des Russischen Reichs. Vol. I. Kayser. Acad. Wiss., St. Petersburg. 504 p.
- Paulson, Dennis R. 1967. Searching for sea serpents. Sea Frontiers 13(4):244-250.
- Peters, Wilhelm K. H., and Giacomo Doria. 1878. Catalogo dei rettili e dei batraci raccolti da O. Beccari, L. M. D'Albertis

- e A. A. Bruijn nella sotto-regione Austro-Malese. Ann. Mus. Civ. Genova 13:323-450.
- Phisalix, Marie. 1912. Modifications que la fonction venimeuse imprime à la tête osseuse et aux dents chez les serpents. Ann. Sci. Natur. Zool. Biol. Anim., ser. 9, 16:161-205.
- 1914. Anatomie comparée de la tête et de l'appareil venimeux chez les serpents. Ibid. 19:1-114.
- Picado, Clodomiro. 1976. Serpientes venenosas de Costa Rica (reprint of 1931 ed.). Editorial Universidad Costa Rica, San Jose. viii + 241 p.
- Pickwell, George V. 1971. Knotting and coiling behavior in the pelagic sea snake *Pelamis platurus* (L.). Copeia 1971(2):348-350.
- 1972. The venomous sea snakes. Fauna 1972(4):17-32.
- , James A. Vick, William H. Shipman, and Marie M. Grenan. 1973. Production, toxicity and preliminary pharmacology of venom from the sea snake, *Pelamis platurus*, with observations on its probable threat to man along Middle America, p. 247-265. In Leonard R. Worthen (ed.), Food-drugs from the sea, Proceedings. Marine Technology Society, Washington, D.C. xix + 396 p.
- Rafinesque, C. S. 1817. Dissertation on water snakes, sea snakes and sea serpents. Amer. Monthly Mag. Crit. Rev. 1:431-435.
- Rosenberg, Herbert I. 1967. Histology, histochemistry, and emptying mechanism of the venom glands of some elapid snakes. J. Morphol. 123(2):133-156.
- Rubinfoff, I., and C. Kropach. 1970. Differential reactions of Atlantic and Pacific predators to sea snakes. Nature 228:1288-1290.
- Russell, Patrick. 1796. An account of Indian serpents collected on the coast of Coromandel. Nicol, London. viii + 91 p.
- Schneider, Johann G. 1799. Historiae amphibiorum naturalis et literariae. Fasc. primus. Frommann, Jena. xiii + 266 p.
- Seba, Albert. 1735. Locupletissimi rerum naturalium thesauri accurata descriptio et iconibus artificiosissimis expressio, per universam physices historiam. Vol. 2. Amstelædami. 154 p.
- Shaw, Charles E. 1961. Snakes of the sea. Zoonooz, San Diego 34(7):3-5.
- 1962. Sea snakes at the San Diego Zoo. Int. Zoo Yearb. 4:49-52.
- Sherborn, C. Davies, and F. A. Jentink. 1895. On the dates of the parts of Siebold's "Fauna Japonica" and Giebel's "Allgemeine zoologie" (first edition). Proc. Zool. Soc. London 1895:149-150.
- Shuntov, V. P. 1963. Morskiye zmei. Priroda (Moscow) 3:103-104.
- Smith, Hobart M., and Edward H. Taylor. 1945. An annotated checklist and key to the snakes of Mexico. U.S. Nat. Mus. Bull. (187):iv + 239.
- Smith, Malcolm. 1920. On sea snakes from the coasts of the Malay Peninsula, Siam and Cochin-China. J. Federated Malay States Mus. 10(1):1-63.
- 1926. Monograph of the sea-snakes (Hydrophiidae). British Museum (Natural History), London. xvii + 130 p.
- Stejneger, Leonhard. 1911. The batrachians and reptiles of Formosa. Proc. U.S. Nat. Mus. 38:91-114.
- Stiles, C. W. 1910. Summaries of six opinions (9, 11, 13, 15, 17, 18) by the International Commission on Zoological Nomenclature. Science 31(787):150-151.
- Stoliczka, F. 1870. Observations on some Indian and Malayan Amphibia and Reptilia. J. Asiatic Soc. Bengal 39(3):159-228.
- Strauch, Alexander. 1873. Die Schlangen des Russischen Reichs in systematischer und zoogeographischer Beziehung. Mem. Acad. Imp. Sci. St.-Peterbourg, ser. 7, 21(4):1-287.
- Taylor, Edward H. 1950. The snakes of Ceylon. Univ. Kansas Sci. Bull. 33, pt. 2(14):519-603.
- Temminck, C. J., and H. Schlegel. 1838. Ophidia, p. 81-96. In Ph. Fr. de Siebold, auctore, Fauna Japonica. C. J. Temminck and H. Schlegel, Reptilia. Müller, Amstelodami. xxi + 144 p.
- Tu, Anthony T. 1976. Investigation of the sea snake, *Pelamis platurus* (Reptilia, Serpentes, Hydrophiidae), on the Pacific coast of Costa Rica, Central America. J. Herpetol. 10 (1):13-18.
- Visser, John. 1967. Color varieties, brood size, and food of South African *Pelamis platurus* (Ophidia: Hydrophiidae). Copeia 1967(1):219.
- Voris, Harold K. 1969. Evolution of the Hydrophiidae with a critique on methods of approach. Ph.D. Thesis, Univ. Chicago. xiv + 429 p.
- 1971. New approaches to character analysis applied to the sea snakes (Hydrophiidae). Syst. Zool. 20(4):442-458.
- 1975. Dermal scale-vertebra relationships in sea snakes (Hydrophiidae). Copeia 1975(4):746-757.
- , C. Kropach, and B. Henderson. 1970. Population biology of *Pelamis*, p. 36. In Alpha Helix research program: 1969-1970. Univ. Calif. San Diego. 59 p.
- , and Bruce C. Jayne. 1976. The costocutaneous muscles in some sea snakes (Reptilia, Serpentes). J. Herpetol. 10(3):175-180.
- Vosmaer, Arnoux. 1774. Beschryving van twee . . . Plaatstaart Slangen. 8 p. Monogr. No. 23. In Natuurkundige beschryving eener uitmuntende verzameling van zeldsaame Gedieren, Bestaande in Oost- en Westindische viervoetige Dieren, Vogelen en Slangen. Meijer, Amsterdam (this constitutes a part of the author's larger Regnum Animale).
- Wagler, Johann. 1830. Natürliches System der Amphibien mit vorangehender Classification der Säugethiere und Vögel. Cotta'schen Buchhandlung, Munich. vi + 354 p.
- Wall, Frank. 1919. A popular treatise on the common Indian snakes. J. Bombay Natur. Hist. Soc. 26:803-810.
- 1921. Ophidia Taprobanica or the snakes of Ceylon. Cottle, Colombo. xxii + 581 p.
- Werner, Franz. 1900. Die Reptilien- und Batrachierfauna des Bismarck-Archipels. Mitt. Zool. Sammlung Mus. Natur. Berlin 1(4):1-132.
- West, G. S. 1895. On the buccal glands and teeth of certain poisonous snakes. Proc. Zool. Soc. London 1895(52):812-826.
- Zeiller, Warren. 1969. Maintenance of the yellowbellied sea-snake, *Pelamis platurus*, in captivity. Copeia 1969(2):407-408.

GEORGE V. PICKWELL, NAVAL OCEAN SYSTEMS CENTER, SAN DIEGO, CALIFORNIA 92152, AND WENDY A. CULOTTA, CALIFORNIA STATE UNIVERSITY AT LONG BEACH, LONG BEACH, CALIFORNIA 90840.

Primary editor for this account, Larry David Wilson.

Published 13 August 1980 and Copyright 1980 by the SOCIETY FOR THE STUDY OF AMPHIBIANS AND REPTILES.